
A Living Landscape Delivering conservation at

a landscape-scale

Our country's wildlife is in trouble.

Between 2002 and 2013, 53% of species in the UK declined, with 41% showing strong or moderate declines (State of Nature, 2016). This includes once common species such as hedgehogs, house sparrows and common toads. Alongside climate change, the main cause is changes in land use, for agriculture and urban development, that have resulted in the loss, fragmentation or inadequate management of habitats.





BETWEEN 2002-2013 SHOWING STRONG OR MODERATE DECLINES

SPECIES IN THE **UK DECLINED**



It is not all bad news.

Our conservation know-how continues to improve and some species populations are recovering, including otters, stone curlews and red kites. Certainly without our protected areas and conservation action to date, the scale of losses would undoubtedly have been much worse.

But the current approach is not enough.

Across the country, and even within our protected sites, the diversity and abundance of wildlife continues to decline and there are ever more pressures facing the environment.

Making Space for Nature

Our response is to make space for nature: moving away from simply trying to conserve what remains and instead restoring and creating habitat at a landscape scale. We call it A Living Landscape.

This approach is based on the concept of 'ecological networks'. At the core of such networks is a collection of high quality sites capable of sustaining species populations. But crucially these core sites need to be connected, so that wildlife is able to move between sites through the wider landscape. And the whole must be robust and resilient, allowing nature to respond to the inevitable changes faced by the environment.

The **Living Landscape** approach must be ambitious, but also realistic: this is not about turning the whole countryside into a nature reserve, but about working alongside other land uses to ensure that we make space for nature.







Size Matters

Larger sites, with their more diverse habitats, will support bigger species populations. Although the size of any population naturally varies in relation to environmental factors, small populations, because they breed less successfully and are unable to generate the numbers required to colonise new sites, are more likely to fluctuate to the point of extinction.

Edae Effects

Compared with the centre of a block of habitat, its edges are likely to be impacted by different climatic conditions, pollution and disturbance arising from adjacent land uses, and different species interactions (including predation). The proportion of a site subject to these edge effects increases the smaller a site becomes, making some or all of that site less suitable for certain species.

Site Proximity

The long term survival of species depends on individuals being able to move between sites and so the closeness of sites to one another is important. The greater the distance between blocks of good quality habitat, the fewer species those blocks will support, since colonisation from other sites is less likely to occur following a decline or extinction.

Meta-Populations

Many species exist not as isolated populations but as sets of populations linked by the movement of individuals between them. Individual populations may decline or even be completely lost, but will re-establish through re-colonisation from nearby sites. If a habitat patch is lost, however, this can impact the whole metapopulation, leading to a decline or even extinction (even if remaining habitat is in good condition).



Better, Bigger, More, Joined

The principles of 'better, bigger, more, and joined', as described by Professor Sir John Lawton in his report Making Space for Nature (2010), provide a valuable framework to guide the effective delivery of **A Living Landscape**.

Better


At the core of any ecological network are existing wildlife habitats, in particular protected sites such as nature reserves, SSSIs and County Wildlife Sites. These support the largest part of our remaining biodiversity and, provided species populations are sufficiently strong, will act as the sources to colonise new areas in the future. Maximising the quality of these core sites 1 is the key first step in delivering A Living Landscape.

The majority of UK habitats are semi-natural: the result of human activity. Maintaining and improving them therefore requires the continuation or re-establishment of conservation management such as grazing or mowing. Even with this on-going management, sites and the wildlife they support will inevitably change in the future, not least due to climate change. However, provided they retain the underlying characteristics which allow them to support a high biodiversity, these sites will remain core to the Living Landscape.





WILDLIFE TRUST increasing the size of core sites creating stepping stone habitats maximising the quality of core sites Creating a Living Landscape buffering habitats to reduce edge effects management practices are as benign as possible 8 ensure land





Bigger

The historic loss and fragmentation of habitats across the country means that surviving patches are generally much smaller – critically so for habitats such as chalk grassland and species-rich wildflower meadows. Increasing the size of core sites through adjacent habitat restoration or creation 2 is required alongside quality improvement. Larger sites will not only support a greater abundance of wildlife, but they can be managed more cost-effectively and naturally.

While making remaining wildlife habitats better and bigger will improve the survival chances of our biodiversity, decades of habitat loss and fragmentation mean that this alone will not be enough to halt further losses. This is where the real step change needs to come.




More

More space needs to be created for wildlife. This requires the establishment of new habitat 3 strategically located according to the needs of the local landscape and its wildlife, to replace at least some of what has been lost. Whatever is created must be as large and of as good a quality as possible.

Joined

The long term survival of wildlife requires that individuals move between patches of core habitat, either to sustain metapopulations or to allow adaptation in light of environmental change, including climate change. Ensuring this movement requires greater connectivity between habitats.



This connectivity may take the form of continuous habitat corridors 4 such as hedgerows between blocks of woodland, or ditches between wetlands. Appropriately managed, these linkages can either support populations in their own right, or can enhance the dispersal of populations.

But direct links are not essential. Creating stepping stone habitats 5 can also act to increase connectivity. While generally smaller than core habitats and so not necessarily capable of supporting sustainable populations in their own right, their location between core sites provides shelter, feeding and resting opportunities for species moving through the countryside.

Maintaining mosaics of mixed land cover **6** can further aid landscape connectivity. The habitat diversity provided by river corridors, networks of ponds and other manmade features such as canals and railway embankments may not match the habitat at the core of an ecological network, but these can all increase the permeability of the countryside, helping species to move across the landscape.

Finally there is a need to reduce the impact on wildlife of our wider land usage. This includes buffering habitats to reduce direct edge effects 7; and ensuring that land management practices are as benign as possible 8 (for example through targeted use of agro-chemicals and reducing other pollution risks).



